Estimating the Size of the MSM Population Using Multiple Methods and Data Sources in Vancouver, British Columbia

A. Rich (1), N.J. Lachowsky (1,2), N. Oliveira (1), G. Colley (1), P. Sereda (1), J. Wong (3), A. Lal (1), J. Jollimore (4), E. A. Roth (5,6), R. S. Hogg (1,7), D. M. Moore (1,2,3)

1. British Columbia Centre for Excellence in HIV/AIDS, Vancouver, BC 2. Faculty of Medicine, University of British Columbia, Vancouver, BC 3. British Columbia Centre for Disease Control, Vancouver, BC 4. Health Initiative for Men (HIM) 5. Department of Anthropology, Faculty of Social Sciences, University of Victoria, Victoria, BC 6. Centre for Addictions Research BC, Victoria, BC 7. Faculty of Health Sciences, Simon Fraser University, Burnaby, BC

Background

- Gay, bisexual and other men who have sex with men (GBMSM) were the first population affected by the HIV epidemic in BC and continues to be the most disproportionately impacted group in the province.
- Lack of a reliable GBMSM population size estimate impedes research and epidemiologic depictions of the HIV epidemic.
- Globally, estimates of GBMSM populations, particularly those in urban centres, are predominantly based on population surveys, capture/recapture estimates, and indirect sampling through serodiagnostic databases. More diverse methodology is necessary for a more accurate understanding of the HIV epidemic.

Methods

We estimated the Metro Vancouver GBMSM population size drawing on four data sources:

- <u>Momentum Health Study</u> cross-sectional survey of GBMSM aged >15 years recruited from February 2012 – February 2014 using respondent-driven sampling (RDS)
- <u>British Columbia Centre for Disease Control (BC CDC) HIV testing data</u> from three sexual health clinics serving GBMSM, February 2012 February 2014
- <u>2011-2012 Canadian Community Health Survey (CCHS)</u> administered by Statistics Canada
- Facebook (social networking website)

Estimates were calculated using the following two methods:

1) Indirect method where *N* is the population estimate, *n* is the number of GBMSM in the group and *p* is the proportion of the Momentum Health Study participants self-reporting such membership. Weights for Momentum estimates were developed using respondents' reported social network size and each corresponding variable.

N = n/p

2) 'Wisdom of The Crowds' (WOTC) method was used to produce an additional point estimate based on the mean of Momentum participants' estimates of the local GBMSM population size.

Results

- **Table 1** presents the range of population estimates.
- CCHS and Facebook data resulted in the lowest population estimates (22,100 and 23,700 respectively), though both sources required disclosure of one's gender and sexual identity.
- HIV testing clinic data and the WOTC method produced estimates (44,300 and 45,800 respectively) nearly double those based on CCHS and Facebook.
- The 2011 Vancouver census male population aged 15+ was 948,010, resulting in GBMSM population estimates ranging from 2.3-4.8%. This estimated range is similar to what the United States (US) Centre for Disease Control (CDC) reports for the US.

Table 1. Population estimates by data source

Data Source & Method	Estimate (N)	n	p	Notes
WOTC	45,800 (IQR: 200-75,000)	-	-	Mean estimate and IQR based on Momentum question: "To the best of your knowledge, how many men who have sex with men, whether they identify as gay or not, do you think live in the Greater Vancouver region?"
Facebook	23,700	Number of Metro Vancouver Facebook profiles specifying gender identity "Male" and preferred partner gender "Interested in men" or " men and women": 18,200	Proportion of Momentum participants reporting a Facebook profile: 76.60 %	Facebook data is based on optional disclosure of gender and sexual identity in user profiles
Sexual Health Clinic HIV Testing Data, adjusted average	44,300	Number of HIV tests for men at 3 clinics serving GBMSM: 5,084	Proportion of Momentum participants reporting HIV testing at any of the 3 GBMSM clinics in the past 2 years: 7.79%	Momentum estimate is adjusted for the average number of HIV tests Momentum participants reported in the past 2 years
Canadian Community Health Survey (CCHS) 2011-2012	22,100	-	<u>-</u>	CCHS data is based on disclosure of sexual identity on government-sponsored, interviewer-administered questionnaire CCHS question: "Do you consider yourself to be: 1) heterosexual (sexual relations with people of the opposite sex); 2) homosexual, that is lesbian or gay (sexual relations with people of your own sex); 3) bisexual (sexual relations with people of both sexes)?"

Discussion

- > Using multiple data sources, our estimates of the Metro Vancouver GBMSM population (22,120-45,800 or 2.3-4.8% of the metropolitan male population) are similar to estimates in other jurisdictions.
- > These findings will support better epidemiologic understanding of the HIV epidemic among GBMSM as well as policy, prevention, and care program decision-making.























